

PYRONOX No. 5

INSECTICIDE

Industrial and Mill Spray

For Rapid Control of Flying and Crawling Insects

Use in FOOD PROCESSING PLANTS, Canneries, Packing Houses, Mills, Elevators, Granaries, Warehouses, Box Cars, Ship's Holds, Breweries, Bottling Plants, Bakeries, Stores, Markets, Theaters, Hospitals, Schools, Hotels, Motels, Restaurants.

ACTIVE INGREDIENTS:

*Piperonyl Butoxide Technical.....	1.50%
Pyrethrins.....	0.30%
INERT INGREDIENTS.....	98.20%
TOTAL	100.00%

*Equivalent to 1.2% (butylcarbityl) (6-propylpiperonyl) ether and to 0.3% of related compounds

KEEP OUT OF REACH OF CHILDREN
CAUTION
Harmful if swallowed.
See side panel for additional precautionary statements.

E.P.A. Reg. No. 45385-30
E.P.A. Est. No. 45385-OH-01

DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal:
Pesticide, spray mixture or rinsate that cannot be used according to label instruction must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal:
Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State or local procedures.

Directions for use continued on back panel

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wear mask or respirator of type recommended by NOISH to give adequate protection against this material. Wear protective clothing. Wash exposed skin with soap and water after handling and before eating. Wash all contaminated clothing with soap and water, separately from other clothing, before reuse.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packaging plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with an acceptable detergent solution and rinsed with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: • Call a physician or Poison Control Center immediately. Gastric Lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING. Vomiting may cause aspiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply directly to water.

PHYSICAL HAZARDS

Do not store near heat or open flame.

Net 1 Gallon / 3.785 Liters

DIRECTIONS, continued

For Maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control **roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, red rust flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites and cadelles:** Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For **silverfish**, spray bookcases. For **ants**, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control **carpet beetles:** Spray edges of carpeting and under carpeting and rugs. Make localized applications to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control **bedbugs:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat as needed.

To control **fleas and brown dog ticks:** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat as needed.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths:** direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of one gallon to 150,000 cubic feet of space.

To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 80,000 cubic feet of space. Contact insects directly whenever possible.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at **1-800-858-7378**



SPECIMEN LABEL